

A NOTE ON A QUESTION OF DIMCA AND GREUEL

PATRICIO ALMIRÓN AND GUILLEM BLANCO

ABSTRACT. In this note we give a positive answer to a question of Dimca and Greuel about the quotient between the Milnor and Tjurina numbers of an isolated plane curve singularity in the cases of one Puiseux pair and semi-quasi-homogeneous singularities.

1. INTRODUCTION

Let $f : (\mathbb{C}^2, \mathbf{0}) \rightarrow (\mathbb{C}, 0)$ with $f(\mathbf{0}) = 0$ be a germ of a morphism defining an isolated plane curve singularity. Associated to any isolated plane curve singularity f one has the Milnor number μ and the Tjurina number τ that are defined as

$$\mu := \dim_{\mathbb{C}} \frac{\mathbb{C}\{x, y\}}{(\partial f / \partial x, \partial f / \partial y)}, \quad \tau := \dim_{\mathbb{C}} \frac{\mathbb{C}\{x, y\}}{(f, \partial f / \partial x, \partial f / \partial y)}.$$

In [3], Dimca and Greuel posed the following question:

Question 1. *Is it true that $\mu/\tau < 4/3$ for any isolated plane curve singularity?*

Furthermore, they show with an example that this bound is asymptotically sharp.

The purpose of this note is to show that Question 1 has a positive answer, using some known results in two cases: the case of one Puiseux pair and for semi-quasi-homogeneous singularities. By a well-known result of Zariski [6], the later case contains the former. However, we decided to include both proofs as the approaches are fundamentally different and may lead to different more general cases of the question. The proof for the first case is based on the results of Delorme [2] and Teissier [5]. For the second case, we use the ideas of Briançon, Granger and Maisonobe [1]. We also show at the end of this note that the bound is also fulfilled by a non-trivial family with two Puiseux pairs studied by Luengo and Pfister [4]. All this gives further evidences for a positive answer of the question in the general case.

Acknowledgments. The authors would like to thank, M. Alberich-Carramiñana, M. González-Villa, A. Melle-Hernández and J. Álvarez-Montaner for the helpful comments and suggestions.

2010 *Mathematics Subject Classification.* Primary 14H20; Secondary 14H50, 32S05.

Key words and phrases. Curve singularities, Tjurina number, Milnor number.

The first author was partially supported by Spanish Ministerio de Ciencia, Innovación y Universidades MTM2016-76868-C2-1-P. The second author was supported by Spanish Ministerio de Ciencia, Innovación y Universidades MTM2015-69135-P and Generalitat de Catalunya 2017SGR-932 projects.

2. ONE PUISEUX PAIR

In this section we will assume that f has a single Puiseux pair (n, m) . We will denote by $\Gamma = \langle n, m \rangle$, $n < m$ with $\gcd(n, m) = 1$ the semigroup of f . Following the ideas of Zariski in [6], Delorme [2] computed the dimension of the generic component $q_{n,m}$ of the moduli space of plane branches with a single Puiseux pair (n, m) .

Theorem 1 ([2, Thm. 32]). *Consider the continued fraction representation $m/n = [h_1, h_2, \dots, h_k]$, with $k \geq 2$, $h_1 > 0$ and $h_2 > 0$. Define, inductively, the following numbers*

$$r_k := 0, \quad t_k := 1, \quad r_{i-1} := r_i + t_i h_i, \quad t_{i-1} := \begin{cases} 0, & \text{if } t_i = 1 \text{ and } r_{i-1} \text{ even,} \\ 1, & \text{otherwise.} \end{cases}$$

Then, the dimension $q_{n,m}$ of the generic component of the moduli space is given by

$$q_{n,m} = \frac{(n-4)(m-4)}{4} + \frac{r_0}{4} + \frac{(2-t_1)(h_1-2)}{2} - \frac{t_1 t_2}{2}.$$

In particular, except for the case $(n, m) = (2, 3)$,

$$(1) \quad \frac{(n-4)(m-4)}{4} \leq q_{n,m} \leq \frac{(n-3)(m-3)}{2}.$$

The bound in the left-hand side of Equation 1 is sharp, consider for instance, the characteristic pair $n = 8, m = 11$. In the Appendix [5] of [6], Teissier, using the monomial curve C^Γ , proves that, in general, the dimension q of the generic component of the moduli space of plane branch with semigroup Γ is given by

$$(2) \quad q = \tau_- - (\mu - \tau_{\min}),$$

where τ_- is the dimension of the miniversal constant semigroup deformation of the monomial curve C^Γ . For one characteristic exponent it is easy to see that

$$\tau_- = \frac{(n-3)(m-3)}{2} + \left\lfloor \frac{m}{n} \right\rfloor - 1,$$

where $\lfloor \cdot \rfloor$ denotes the integer part. In this case, the Milnor number is $\mu = (n-1)(m-1)$. Combining the lower bound in Equation 1 and Equation 2 one obtains the following lower bound for τ_{\min}

$$(3) \quad \frac{(n-4)(m-4)}{4} + (n-1)(m-1) - \frac{(n-3)(m-3)}{2} - \frac{m}{n} + 1 \leq \tau_{\min}.$$

except for the case $(n, m) = (2, 3)$.

Proposition 1. *For any plane branch with one characteristic exponent, $\mu/\tau < 4/3$.*

Proof. It is sufficient to proof the inequality for the τ_{\min} of each characteristic pair (n, m) . Dividing μ by the expression in Equation 3 and rewriting

$$(4) \quad \frac{\mu}{\tau} \leq \frac{\mu}{\tau_{\min}} \leq \frac{4n(n-1)(m-1)}{3n^2m - 2n^2 - 2nm + 6n - 4m},$$

assuming always that $(n, m) \neq (2, 3)$, $n < m$. The upper bound in Equation 4 is strictly smaller than $4/3$ if and only if $0 < m(n-4) + n(n+3)$. Therefore, the result holds if $n \geq 4$. The cases $n = 2$ and $n = 3$ follow from computing the τ_{\min} using Theorem 1.

Indeed, let $n = 2$ and $m = 2h_1 + 1$, $h_1 > 1$ so the continued fraction representation is $m/n = [h_1, 2]$. Then, $r_0 = 2$, $t_1 = 0$, $t_2 = 1$ and $q_{2,m} = h_1 - m/2 - 1/2 = 0$. Analogously, if $n = 3$, then $m = 3h_1 + 1$ or $m = 3h_1 + 2$; the continued fractions are either $m/n = [h_1, 3]$ or $m/n = [h_1, 1, 2]$. Then, $r_0 = 3 + h$ or $r_0 = 2 + h$, $t_2 = 1$ or $t_2 = 0$, respectively, and $t_1 = 1$ in either case. Consequently, in both cases, $q_{3,3h_1+1} = -m/4 + 3h_1/4 + 1/4 = 0$ and $q_{3,3h_1+2} = -m/4 + 3h_1/4 + 1/2 = 0$. Finally, since $\tau_- = 0$ if $n = 2$ and $\tau_- = h_1 - 1$ if $n = 3$,

$$\frac{\mu}{\tau_{\min}} = 1 < \frac{4}{3}, \quad \frac{\mu}{\tau_{\min}} < \frac{6m-6}{5m-3} < \frac{6}{5} < \frac{4}{3},$$

for $n = 2, m \geq 3$ and $n = 3, m \geq 4$, respectively. \square

3. SEMI-QUASI-HOMOGENEOUS SINGULARITIES

We assume now that f is a semi-quasi-homogeneous singularity with weights $w = (n, m)$ such that $\gcd(n, m) \geq 1$ and $n, m \geq 2$. This means that $f = f_0 + g$ is a deformation of the initial term $f_0 = y^n - x^m$ such that $\deg_w(f_0) < \deg_w(g)$. In [1], Briançon, Granger and Maisonobe, using the technique of escaliers, give recursive formulas to compute the τ_{\min} of this type of singularities. Their main result is the following:

Theorem 2 ([1, §I.6]). *For semi-quasi-homogeneous singularities with initial term $y^n - x^m$,*

$$\tau_{\min} = (m-1)(n-1) - \sigma(m, n).$$

The number $\sigma(a, b)$ is defined recursively for any non-negative integers a, b as follows. If $a, b \leq 2$ then $\sigma(a, b) := 0$. Otherwise, we can express $a = bq + r$, $0 \leq r < b$, $q \geq 1$. For the cases $r = 0, 1, b-1, b/2$ there are closed formulas for $\sigma(a, b)$ denoted by $\Sigma_0, \Sigma_1, \Sigma_{b-1}, \Sigma_{b/2}$, see Table 1 in [1]. If none of the above cases hold, define recursively, see Tables 2 and 3 in [1], a finite sequence $(a_0, b_0), (a_1, b_1), \dots, (a_k, b_k)$ with $(a_0, b_0) = (m, n)$, $\sigma(a_k, b_k)$ is in one of the previous cases, and for $i = 0, \dots, k-1$:

(A) If $\gcd(a_i, b_i) = 1$, we can find $ub_i - va_i = 1$ with $2 \leq u < a_i$. Letting $\gamma := [\frac{a_i}{u}]$, we have two subcases:

(AE) If γ is even, define $a_{i+1} = a_i - \gamma u$, $b_{i+1} = b_i - \gamma v$, then

$$\sigma(a_i, b_i) := \frac{(a_i - 2)(b_i - 2)}{4} - \frac{(a_{i+1} - 2)(b_{i+1} - 2)}{4} - \frac{\gamma}{4} + \sigma(a_{i+1}, b_{i+1}).$$

(AO) If γ is odd, define $a_{i+1} = (\gamma + 1)u - a_i$, $b_{i+1} = (\gamma + 1)v - b_i$, and

$$\sigma(a_i, b_i) := \frac{(a_i - 2)(b_i - 2)}{4} - \frac{(a_{i+1} - 2)(b_{i+1} - 2)}{4} - \frac{\gamma + 1}{4} + \sigma(a_{i+1}, b_{i+1}).$$

(B) Otherwise, $a_i = \alpha a'$, $b_i = \alpha b'$ with $\alpha \geq 2$, $\gcd(a', b') = 1$, and we can find a Bezout's identity $ub' - va' = 1$ with $1 \leq u < a'$. We have again two subcases:

(BP) If α is even,

$$\sigma(a_i, b_i) := \frac{(a_i - 2)(b_i - 2)}{4} - \frac{\alpha}{2}.$$

(BO) If α is odd, define $a_{i+1} = |a' - 2u|$ and $b_{i+1} = |b' - 2v|$, and

$$\sigma(a_i, b_i) := \frac{(a_i - 2)(b_i - 2)}{4} - \frac{\alpha}{2} - \frac{(a_{i+1} - 2)(b_{i+1} - 2)}{4} + \sigma(a_{i+1}, b_{i+1}).$$

Proposition 2. *For any semi-quasi-homogeneous singularities with initial term $y^n - x^m$,*

$$\mu/\tau < 4/3.$$

Proof. Observe that in the recursive cases (A) and (BO),

$$\sigma(a, b) \leq \frac{(a-2)(b-2)}{4} - \frac{(a_k-2)(b_k-2)}{4} + \sigma(a_k, b_k),$$

where $\sigma(a_k, b_k)$ is either zero or has a closed form. Notice also that $a_i b_{i+1} > b_i a_{i+1}$ for all $i = 0, \dots, k-1$. From these observations, one can deduce that, in general,

$$(n-1)(m-1) - \frac{(m-2)(n-2)}{4} - \kappa(n, m) \leq \tau_{\min},$$

where $\kappa(n, m) = m/4n$ if $\sigma(a_k, b_k)$ is $\Sigma_0, \Sigma_1, \Sigma_{b-1}$ with b odd, $\kappa(n, m) = 5/4$ if $\sigma(a_k, b_k)$ is $\Sigma_0, \Sigma_1, \Sigma_{b-1}$ with b even or $\Sigma_{b/2}$ with $b/2$ odd, and $\kappa(n, m) = 0$ if $\sigma(a_k, b_k)$ is $\Sigma_{b/2}$ with $b/2$ even or in the case (BP). In any case,

$$\frac{\mu}{\tau} \leq \frac{\mu}{\tau_{\min}} \leq \frac{4(n-1)(m-1)}{3nm - 2n - 2m - 4\kappa(n, m)},$$

which is bounded by $4/3$ if and only if $n+m+\kappa(n, m) > 3$, which is true for $n, m \geq 2$. \square

4. A FAMILY WITH TWO PUISEUX PAIRS

In [4], Luengo and Pfister study the family of irreducible plane curve singularities with semigroup $\langle 2p, 2q, 2pq + d \rangle$ such that $\gcd(p, q) = 1, p < q$ and d odd. The Milnor number of this family equals

$$\mu = (2p-1)(2q-1) + d.$$

Studying the kernel of the Kodaira-Spencer map, they prove, see [4, pg. 259], that τ is constant in each equisingularity class and equals,

$$\tau = \mu - (p-1)(q-1).$$

One can easily check that $\mu/\tau < 4/3$ for all the semigroups of the family.

REFERENCES

- [1] J. Briançon, M. Granger, Ph. Maisonobe, *Le nombre de modules du germe de courbe plane $x^a + y^b = 0$* , Math. Ann. **279** (1988), 535-551.
- [2] C. Delorme, *Sur les modules des singularités des courbes planes*, Bull. Soc. Math. France **106** (1978), 417-446.
- [3] A. Dimca, G.-M. Greuel, *On 1-forms on isolated complete intersection on curve singularities*, arxiv.org/pdf/1709.01451.pdf, (2017).
- [4] I. Luengo, G. Pfister, *Normal forms and moduli spaces of curve singularities with semigroup $\langle 2p, 2q, 2pq + d \rangle$* , Compos. Math. **76** (1990), no.1-2, 247-264.
- [5] B. Teissier, *Appendix*, in [6], 1986.
- [6] O. Zariski, *Le problème des modules pour les branches planes*, Hermann, Paris, 1986.

DEPARTAMENT DE MATEMÀTIQUES, UNIV. POLITÈCNICA DE CATALUNYA, AV. DIAGONAL 647, BARCELONA 08028, SPAIN.

E-mail address: Guillem.Blanco@upc.edu

DEPARTAMENTO DE ÁLGEBRA, GEOMETRÍA Y TOPOLOGÍA, FACULTAD DE CIENCIAS MATEMÁTICAS,, UNIVERSIDAD COMPLUTENSE DE MADRID, 28040, MADRID, SPAIN.

E-mail address: palmiron@ucm.es